

Ed Clyde

Brooke
sis to Sophia
Brooke & Jane
Hatch

Jane Hatch
Jo's wife

Mary Willis
faced east
Photography

UTAH INFLUENZA IMMUNIZATION PROJECT

INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at

Frank Epperson
sang

J. E. McMullen

F = Henry

M Jeannette
Murdock

~~from~~

Henry Moulton

taught school

UTAH INFLUENZA IMMUNIZATION PROJECT

INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at

John E Medley
Trucker

Lottie & Fred
Hiles

John

addy

Sissy

Albert

Sewey

UTAH INFLUENZA IMMUNIZATION PROJECT

INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at